

Consistent discretizations for classical and quantum general relativity

Rodolfo Gambini¹, and Jorge Pullin²

1. Instituto de Física, Facultad de Ciencias, Universidad de la República, Iguaú 4225, CP 11400 Montevideo, Uruguay

2. Department of Physics and Astronomy, Louisiana State University, 202 Nicholson Hall, Baton Rouge, LA 70803-4001
(August 23th 2001)

We consider discretizations of the Einstein action of general relativity such that the resulting discrete equations of motion form a consistent constrained system. Upon “spin foam” quantization of the system, consistency allows a natural way of recovering the correct semi-classical theory. A consistent set of approximations to the Einstein equations could also have implications for numerical relativity and for the construction of approximate classical observables for the theory.

There has been significant growth in interest in computing the path integral for general relativity [1] due to the introduction of mathematical tools [2] that allow to control the space of quantum states of the gravitational field. The approach is generically known as “spin foam” approach to quantum gravity. It consists in postulating discrete approximations to the path integral of general relativity in terms of a simplicial decomposition of space-time. There has been steady progress in showing that, at least for a given discretization, there is a chance that the discrete path integral can be finitely computed, even in the Lorentzian case [3]. Part of the motivation for this approach comes from the success in spin foam approaches to quantize topological field theories [4]. However, there is a significant difference between topological field theories and general relativity. Topological field theories have a finite number of degrees of freedom, whereas general relativity is a genuine field theory with local degrees of freedom. This has important implications at the time of considering the spin foam approach. In topological field theories, one only needs to consider a discretization on a given simplicial decomposition, since the results are independent of the choice of simplicial decomposition of the manifold. In general relativity this is not likely to be the case. One therefore is left with an approximation to the theory that is discretization dependent. The hope is that in the limit in which one infinitely refines the triangulation one will recover the continuum theory through some kind of averaging or limiting procedure. This is problematic. Since general relativity is not perturbatively renormalizable, there is no guiding principle that ensures that the limit will be well defined. For instance, in Yang-Mills theories on the lattice, one can write infinitely many discretizations of the dynamics that are all different but become the same due to the phase transition of the theory that defines the continuum limit. No such procedure is known for gravity. In fact, a point of view advocated some times in the gravitational case is that one should not take the continuum limit since the theory has a natural cutoff length given by the Planck length. This point of view suggests that quantum gravity will be inherently a discrete theory (see for instance the discussion in [3]). Similar comments apply to the canonical quantization of general relativity. In this case, proposals for the Hamiltonian constraint have been made [5], that are based upon discretizations of the theory. Different discretizations appear to lead to different theories [6], which may or may not yield the same semiclassical limits, which are currently beginning to be studied [7]

The above comments suggest that either due to fundamental reasons or due to practical ones, one will be dealing with discretizations of general relativity at the time of quantizing the theory. Discretizations are also useful in a completely different area of physics: numerical simulations of the Einstein equations. Again, one is really interested in the limit in which the discretization is infinitely refined, but in practice one handles a discrete theory when performing a numerical simulation.

A striking element of most discretizations of general relativity currently used is that in and of themselves, they define *inconsistent* theories. For instance, it would be highly desirable if the spin foam models of the path integral stemmed from the quantization of discretizations of the classical action. However, it is quite non-trivial to obtain discretizations of the action of general relativity such that the classical equations of motion stemming from the discrete action have non-trivial solutions and reproduce in the limit the continuum theory at the level of the equations of motion (for a discussion in the context of Regge Calculus see [8]). In the case of numerical relativity, the set of discrete equations that represent the Einstein evolution equations are usually inconsistent with the discrete equations that represent the constraints. It is a bit surprising that these significant difficulties have not attracted attention in the past, but the general attitude towards discretizations has been that whatever inconsistencies one has will disappear in the continuum limit and are therefore tolerable. However, if the continuum limit is not achieved (either by fundamental or practical reasons as we discussed above), we believe these inconsistencies should be taken seriously.

The inconsistencies might appear surprising at first, but they are not. There is a body of literature dealing with the space-time discretization of dynamical systems [9], and even for elementary mechanical systems it is difficult to achieve consistency. As an example, let us consider a simple mechanical system, a particle in a potential, and let us take time to be discrete. The action then reads $S = \sum_{i=1}^N p_i(q_{i+1} - q_i) + [p_i^2/2 + V(q_i)] \Delta t$. If one works out the discrete equations, one gets $q_{i+1} - q_i = p_i$, $p_i - p_{i-1} = -V'(q_i)$. Notice the asymmetry in the left members. This time asymmetry, in turn, implies that energy is not conserved, as can be checked through a straightforward calculation.

This problem also implies that if one adds constraints that are not holonomic (i.e. they mix p_i and q_i) the constraints are not preserved in evolution. Most of the literature up to now on “discrete mechanics” has considered only systems with holonomic constraints (see [10] for references). Such systems therefore do not include general relativity.

In this paper we would like to suggest two new proposals for constructing consistent discretizations of general relativity. The first one will be in the context of discretizing equations of motion in a consistent way, like one would do in numerical relativity. Here what we will seek to find is discrete systems of evolution equations that are compatible with the discrete constraints. The second will be in discretizing an action like is done in spin foam quantum gravity. Here what we will attempt to do is to find a discretization of the classical action that yields discrete equations of motion that are a good approximation to the continuum equations of motion and that admit solutions that approximate the classical solutions. We will argue how this will help in constructing the semi-classical limit of the theory.

Discretizing the Einstein equations. Let us start by considering how to discretize consistently the equations of general relativity. To motivate what we will propose, we would like first to work out a simpler example, closely related to one solved by T.D. Lee [11]. We will see that the technique introduced naturally suggests what to do in the case of general relativity. The example is a one dimensional mechanical system. The discrete action for the system is $S = \sum_n p_n(q_{n+1} - q_n) - H(p_n, q_n)(t_{n+1} - t_n)$. The novelty in T.D. Lee’s approach is to consider the t_n as canonical variables, the theory being “discretely parameterized” by n . Variation with respect to t_n shows that $H_n = H_{n-1}$ and therefore the Hamiltonian is automatically conserved. One has three equations of motion, given by variations with respect to t_n, q_n, p_n . The equations will determine t_n, p_n, q_n and therefore the time intervals are determined by the evolution. Let us now consider the same construction for a constrained system. Consider the same system with a constant energy E_0 . The action will read $S = \sum_n p_n(q_{n+1} - q_n) - \lambda_n(H(p_n, q_n) - E_0)$. We have incorporated into the Lagrange multiplier λ_n the time interval. We now have an equation of motion with respect to λ_n which states that the energy is constant and equations of motion with respect to q_n, p_n that will involve the Lagrange multiplier. Solving the system shows that the Lagrange multiplier will be determined by the evolution. Is time reparametrization lost? It is not, since we had absorbed the time interval into the definition of the multiplier, we can always choose a different time interval in the theory. This example suggests what we can do in general relativity. Our proposal consists in taking the $3 + 1$ decomposition of the Einstein equations discretized on a grid, like it is normally done in numerical relativity. One will get discrete time evolution equations for the extrinsic curvature and the three metric. The right hand sides of these equations will involve the (discretized) lapse and shift. We then add to these equations the four discrete constraint equations evaluated at time $n + 1$. The resulting system should be solved for the metric, extrinsic curvature, lapse and shift, in such a way that both evolution and constraint equations are solved. One is then left with a consistent system, at the expense of fixing the lapse and shift. It might appear paradoxical at first that the lapse and shift are fixed, but it is not. Since discrete theories do not have diffeomorphism invariance, in general one has to fix the lapse and shift for the theories to be consistent. This was observed in the context of Regge Calculus by Friedmann and Jack [12]. However, it still seems strange that one is trying to approximate a continuum theory, which has arbitrary lapse and shifts, with a discrete theory that has fixed ones. The answer lies in the fact that when one discretizes a theory there is implicit a mapping between space-time and the discrete lattice. Such mapping is arbitrary and restores the reparametrization freedom of general relativity. It is analogous to what happened in the case of the harmonic oscillator, where what was really fixed was not the Lagrange multiplier, but its value times the discretized time interval. It is evident that what we have up to now is just a proposal, and further exploration in practical numerical implementation will be needed. In particular, although the system constructed solves exactly the discrete constraints (and therefore approximates well the continuum constraints forever in time) our experience with model systems shows that since one needs to solve non-linear equations for the Lagrange multipliers, there are situations where one might only find complex solutions, which will render the scheme useless. Studies applied to one dimensional systems are currently under way. After completion of this manuscript we became aware of work of Van Putten along similar lines [13], who appears to have tested a closely related proposal in the context of Gowdy waves.

Discretizing the action of general relativity and quantization

In the spin foam approach to quantum gravity, a certain discretization of the path integral is proposed. This is not necessarily achieved by a unique prescription. In general what people have in mind would be to write the action of general relativity using connection-type variables, discretized on a given simplicial decomposition. One then performs the functional integrals on the connections and is left with an expression that is given by a sum over “spin foams”, that is, an assignment of group-valued labels to a graph. Due to the discretization, the functional integrals become ordinary integrals on the group (typically $SU(2)$). This “procedure” is loosely followed. Usually, certain expressions for the final discretization are chosen a priori based on certain desirable criteria, they are not necessarily obtained strictly through a discretization of the classical action. As we mentioned in the introduction, the resulting quantum theories are discretization-dependent and it is not clear in what sense they are connected with classical general relativity. It is expected that in some limiting procedure one should recover the semi-classical theory. There is however no guiding principle to take this limit.

Our point of view is that it would be highly desirable to construct spin foam models starting with a consistent

discretization of the action of general relativity, in the sense that before quantization, one has a discrete classical action that yields discrete equations of motion that are consistent and with solutions that in the continuum limit go to those of general relativity. Even with this scenario, one should be aware that discretized theories in general have more solutions than the continuum theory. These solutions have “spikes” or “high frequency” oscillations and do not converge when one refines the discretization to solutions of the continuum theory.

The strategy for computing the functional integral will then be to choose a consistent discretization and compute the transition amplitude for it; this calculation is well defined since one is dealing with a system with a finite number of degrees of freedom and a discrete time. Therefore the transition amplitude is given by an ordinary integral. If we were to stop here and try to use the transition amplitude as an approximation to the path integral of the continuum theory, we would quickly discover that one has a bad approximation. Since the discrete theory contains a large number of spurious solutions, the resulting transition amplitude will in general fail to capture the behavior one associates with the continuum theory. Moreover, the result will be discretization-dependent. This is the typical situation that is being confronted by current approaches to spin-foam quantization. In our approach we have a clear advantage. Since the discretization constructed is consistent, one is ensured that the discrete theory will contain a solution that approximates better and better the continuum solution as one refines the discretization. It does, however, contain many other spurious solutions. To get rid of these we propose the following construction. Consider the discrete theory on a simplicial decomposition of the manifold with a finite number of simplices. We then average the resulting transition amplitude over all possible discretizations with up to the given number of simplices. This is a finite sum, so it is well defined. Each discretization will therefore contain a transition amplitude that approximates the one of the continuum theory and many others that are discretization dependent. The averaging procedure will therefore suppress the discretization-dependent amplitudes, since one in general has only one of each per discretization, and will be dominated by the amplitudes that approximate those of the continuum theory. One can then repeat this averaging with more simplices. As one increases the number of simplices, given that the discretization is consistent, one ensures that the transition amplitude will model better and better that of the continuum theory. What would have happened if one had picked an inconsistent discretization? In such case, the various approximations correspond to theories that are not related to the classical theory in any obvious way. Although one can perform a path integral quantization using an action that yields inconsistent equations of motion, we cannot see a sense in which summing a large number of such contributions will conspire to yield an amplitude related to classical general relativity, unless one could find a guiding principle that could suggest this should happen (e.g. renormalizability).

Let us discuss a consistent discretization suitable for spin foam quantization. The discretization has been proposed by Reisenberger [14]. The new elements in this paper is to indicate that indeed the discretization is consistent at least for simple discretizations. Although the discretization of Reisenberger was introduced for Euclidean gravity, it can be generalized to the Lorentzian case. The functional integrals in that case are problematic since the group is non-compact, but the recent results of [3] suggest that they could be handled through a certain regularization. The model is based on the Plebanski [15] action of general relativity,

$$S = \int \Sigma_i \wedge F^i - \frac{1}{2} \phi^{ij} \Sigma_i \wedge \Sigma_j \quad (1)$$

where ϕ^{ij} is a symmetric traceless matrix, F^i and Σ_i are $SU(2)$ valued two forms. F^i is the curvature of an $SU(2)$ connection. Variation with respect to ϕ^{ij} gives a constraint equation for the Σ_i that implies (at least for non-degenerate solutions) that it is given by a skew product of one-forms $\Sigma^i = e_j \wedge e_k \epsilon^{ijk}$. If one substitutes back in the action, one recognizes the Samuel [16] action for general relativity. One now considers a simplicial decomposition of the manifold Δ . Each 4-simplex ν is bounded by five tetrahedra τ , intersecting at ten faces σ . One can then introduce the “derived” (or “dual”) complex, formed by wedges determined by the following four points: the simplex center C_ν , the centers of two adjacent tetrahedra C_τ , and the centers of the faces C_σ . Each wedge s is bounded by a polygonal line starting at the center of the four simplex, going over to the center of one of the tetrahedra, continuing to the center of a face, over to the center of the other tetrahedra adjoined by that face and returns to the center of the 4-simplex. There is therefore one wedge “dual” to each face. To introduce the discrete variables, the connection is represented by the holonomy along one of the sides of the wedge, the curvature by taking the trace of the product of a closed holonomy along the wedge times a generator of $SU(2)$, which we call θ^i . Each wedge has associated with it an object e_{si} that represents the two form Σ_i . The matrix ϕ^{ij} is assumed based at the center of the 4-simplex, so we denote it ϕ_ν^{ij} . The discrete action is given by [14],

$$S_\Delta = \sum_{\nu \in \Delta} \left[\sum_{s \in \nu} e_{si} \theta^i - \frac{1}{60} \phi_\nu^{ij} \sum_{s, \bar{s} \in \nu} e_{si} e_{\bar{s}j} \text{sgn}(s, \bar{s}) \right] \quad (2)$$

where $\text{sgn}(s, \bar{s})$ is non-vanishing if the two wedges share at most one vertex, and is ± 1 depending on the orientation of the wedges.

A wedge has two types of links, the ones that start from the center of the simplices (“interior links”) and the ones that go from the center of the tetrahedra to the center of the face (“boundary links”). The discrete equations of motion are obtained by varying with respect to the holonomies associated with the two types of links, with respect to ϕ_{ν}^{ij} and with respect to the e_{si} . Varying with respect to the interior links one gets a sum over all the faces of the quantity e_{is} associated with the wedge dual to the face, parallel-transported to the center of the simplex. This sum has to vanish and corresponds to a discretization of Gauss’ law. Varying with respect to the boundary links one gets a holonomy around the wedge with the insertion of an e_{is} with s the dual of the wedge. The equation states that this quantity is the same for all wedges associated with simplices that share the face σ . Varying with respect to ϕ^{ij} one gets a constraint that corresponds in the continuum to the two form Σ being decomposable into the wedge product of two one-forms. Finally, varying with respect to e_{is} one gets an expression that links the “curvature” θ to the multipliers ϕ^{ij} and the field e_{si} . In the continuum this would correspond to relating the curvature with the Weyl tensor. As we have suggested, all the equations can be proved in detail to have the correct continuum limits [14].

Is the resulting theory consistent? In order to establish this in general one would need a full canonical analysis, which is not available at present. We have however been able to establish that the equations of motion are consistent for particular discretizations. As an example let us consider a single simplex. To count variables, we start by choosing a gauge in which the only non-trivial holonomies are associated to the boundary links that are outgoing from the face dual to the wedge on which the link lives. There are 10 such holonomies, and in terms of them we can compute all holonomies along wedges. To see that this gauge exists, one first chooses a gauge transformation at the centers of tetrahedra that makes the internal holonomies identity. One then chooses a gauge transformation in the center of faces in such a way that the incoming holonomies are trivial (one could alternatively choose the outgoing ones). We therefore have 30 variables associated with the holonomies (for $SU(2)$ each holonomy can be parameterized by three real variables). We also have the five ϕ^{ij} ’s and 30 e_{si} ’s, for a total of 65 variables. Counting equations, the Gauss law gives 15 identities (three per each of the five faces). The second set of equations is trivial, since there is only one simplex. The third set of equations gives five conditions and the last set corresponds to 30 conditions. So we have a total of 50 equations for the 65 variables. The rest corresponds to “boundary data” (the boundary of the tetrahedra can be viewed as a sandwich-boundary value problem) that are freely specifiable. Similar countings can be performed for more complex simplicial decompositions. We have failed to find one that leads to inconsistencies.

We therefore see that we have two different discretizations of classical general relativity that are consistent, one more geared towards a numerical implementation and one suited for spin foam quantization. We believe that the use of consistent discretizations will prove fruitful in producing theories with the correct continuum limits, both at a classical and quantum level. Having a consistent discrete theory allows to perform studies of a concrete, definitive nature (for instance computing observables), providing a laboratory for testing the validity of the approximations constructed.

We wish to thank Doug Arnold and Richard Price for comments. This work was supported by grants NSF-PHY090091, NSF-INT-9811610.

-
- [1] See J. C. Baez, Lect. Notes Phys. **543**, 25 (2000) for a recent review and references.
 - [2] A. Ashtekar and J. Lewandowski, J. Geom. Phys. **17**, 191 (1995).
 - [3] L. Crane, A. Perez and C. Rovelli, gr-qc/0104057.
 - [4] H. Ooguri, Nucl. Phys. **382**, 276 (1992).
 - [5] T. Thiemann, Class. Quan. Grav. **15** 839; 875; 1207; 1249; 1281; 1463 (1998); C. Di Bartolo, R. Gambini, J. Griego, J. Pullin, Phys. Rev. Lett. **84** 2314 (2000); Class. Quan. Grav. **17**, 3211; 3239 (2000).
 - [6] M. Gaul and C. Rovelli, Class. Quant. Grav. **18**, 1593 (2001)
 - [7] H. Sahlmann, T. Thiemann and O. Winkler, Nucl. Phys. B **606**, 401 (2001) and references therein.
 - [8] L. C. Brewin and A. P. Gentle, Class. Quant. Grav. **18**, 517 (2001)
 - [9] See G. Jaroszkiewicz and K. Norton, J. Phys. AA **30**, 3115 (1997) and references therein.
 - [10] J. Wendlandt, J. Marsden, Physica **D106**, 223 (1997).
 - [11] T. D. Lee, in “How far are we from the gauge forces” Antonino Zichichi, ed. Plenum Press, (1985)
 - [12] J. L. Friedman and I. Jack, J. Math. Phys. **27**, 2973 (1986).
 - [13] M. Van Putten, “Past and future gauge in numerical relativity” gr-qc/0108056.
 - [14] M. P. Reisenberger, Class. Quant. Grav. **14**, 1753 (1997)
 - [15] J. Plebanski, J. Math. Phys. **18**, 2511 (1977); R. Capovilla, J. Dell, T. Jacobson, Phys. Rev. Lett. **63**, 2325 (1989).
 - [16] J. Samuel, Pramana **28**, L429 (1987).